SmoothHD FOR SINGLE CONTRAST GI FLUOROSCOPY

BARIUM SULFATE SUSPENSION
105% w/v

SmoothHD barium sulfate oral suspension

- Indicated for use in fluoroscopy in single contrast upper and lower gastrointestinal tract.
- 450mL high-density barium suspension formulation for adults and pediatric patients 12 years and older.

Preparation

1. **Shake bottle vigorously for 30 seconds.**

2. **Wait for a minimum of 5 minutes** (allows air bubbles to float out of suspension, improving image quality).

Upper GI
Blend SmoothHD 105% Mixture + Water

a). **Mix Ratio:**
150 mL SmoothHD + 150 mL water

b). **Stir Thoroughly:**



UGI & Small Bowel Follow Through
Prepare 3 Cups of This Mixture

Quantity Required:
Prepare 3 cups of of SmoothHD and water mixture.
Use 1 container (450 mL) of SmoothHD.

Post-Exam Instructions for Patient

Hydration: Encourage plenty of fluids for 48 hrs.

Diet: Patient may eat and drink as desired post-exam.

Safety Information

For oral administration only. This product should not be used in patients with known or suspected perforation of the GI tract, known obstruction of the GI tract, conditions associated with high risk of GI perforation or aspiration, or known hypersensitivity to barium sulfate or any of the excipients in SmoothHD. Intra-abdominal leakage may occur in conditions such as GI fistula, ulcer, inflammatory bowel disease, appendicitis or diverticulitis, severe stenosis or obstructing lesions of the GI tract. Caution is recommended in patients with history of food aspiration and known swallowing disorders. Patients should maintain adequate hydration in days following a barium sulfate procedure to avoid obstruction or impaction. Please consult full Prescribing Information for SmoothHD online at: https://voyageurpharmaceuticals.ca/smooth-hd

VOYAGEUR
FROM THE EARTH TO THE BOTTLE®

These guidelines are suggested uses only final discretion always rests with the attending radiologist.

VoyageurPharmaceuticals.ca + 1-877-398-6866 info@vpharma.ca