SmoothLD FOR SINGLE CONTRAST GI FLUOROSCOPY

BARIUM SULFATE SUSPENSION 60% w/v

SmoothLD barium sulfate suspension 60% w/v

Indicated for use as a low-density (60% w/v) barium suspension solution, SmoothLD is tailored for single contrast radiographic examinations of the upper GI tract and small bowel.

For both adults and children.

Preparation

1. **Shake Well:**

a). **Vigorously shake SmoothLD bottle for 30 seconds before use.**



b). **Shake again just before each patient swallow.**

Shake again if product has been standing for over 60 minutes before administration.

2. **Dosage Form**: Ready-to-use liquid suspension at 60% concentration. Do not dilute or modify.

Dosing & Administration

SmoothLD is ready to use

Post-Exam Instructions for Patient

Hydration: Hydrate well for 48 hrs to prevent constipation.

Diet: Resume normal eating and drinking as tolerated.

Safety Information

For oral administration only. This product should not be used in patients with known or suspected perforation of the GI tract, known obstruction of the GI tract, conditions associated with high risk of GI perforation or aspiration, or known hypersensitivity to barium sulfate or any of the excipients in SmoothLD. Intra-abdominal leakage may occur in conditions such as GI fistula, ulcer, inflammatory bowel disease, appendicitis or diverticulitis, severe stenosis or obstructing lesions of the GI tract. Caution is recommended in patients with history of food aspiration and known swallowing disorders. Patients should maintain adequate hydration in days following a barium sulfate procedure to avoid obstruction or impaction. Please consult full Prescribing Information for : https://voyageurpharmaceuticals.ca/smooth-ld

VOYAGEUR
FROM THE EARTH TO THE BOTTLE®

These guidelines are suggested uses only final discretion always rests with the attending radiologist.

VoyageurPharmaceuticals.ca + 1-877-398-6866 info@vpharma.ca